



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

NOTES AND ERRATA: VOLUMES 1, 2, 3

VOLUME 1

F. R. MOULTON: *On a class of particular solutions . . .*

P. 28, formula (26).	For	$x - \frac{1}{\sqrt{3}}$ $\left\{ \left(x - \frac{1}{\sqrt{3}} \right) \right\}^{\frac{2}{3}}$	read	$x - \frac{1}{\sqrt{3}}$ $\left\{ \left(x - \frac{1}{\sqrt{3}} \right)^2 \right\}^{\frac{2}{3}}$
" 1. 6 up.	"	1.22	"	1.18 .
" 1. "	"	-.24	"	-.46 .
" 1. 5 up.	"	-.93	"	-.77 .

VOLUME 2

L. E. DICKSON: *Canonical forms of quaternary . . .*

P. 107, l. 1.	For	ξ'	read	ξ'_1 .
P. 109, l. 7 up.	"	chose	"	choose.
P. 110, l. 4.	"	$L_{11} T_{1-1}, L_{2\mu}$	"	$L_{11} T_{1-1} L_{2\mu}$.
P. 113, l. 22.	"	determines δ_{12}	"	determines β_{12} .
P. 121, l. 2 of § 15.	"	=	"	+
P. 125, middle.	The number (33) refers only to the first of the two equations.			

G. A. MILLER: *Determination of all the groups of order p^m . . .*

P. 263, l. 5.	For	t^i	read	t_7 .
" 1. 10.	"	$p > 3$	"	$n = 2, 3, \dots, p - 2$.
" 1. 11.	Read	$(t_7^{-1} t_6 t_7)^{-1} P_2^n (t_7^{-1} t_6 t_7) = P_3 P_2^n = P_3^n P_2^n = t_6^{-n} P_2^n t_6^n$.		
P. 271, l. 4.	For	$0, p^4 - 1$	read	$p^2 - 1, p^2(p^2 - 1)$.
Pp. 262, 263.	From the second and third corrections it follows that the			

$p(p-1)$ subgroups of order p , mentioned in the second line from the bottom of p. 262, form two equal conjugate sets, and that the non-invariant subgroups of order p^2 and type $(1, 1)$ contained in I are conjugate under I in sets of $(p-1)/2$ instead of forming a single conjugate set, as is stated in line 18 on p. 263. There are, therefore, eight groups of order p^m ($p > 2$) which are non-abelian and include the abelian group of type $(m-2, 1)$; i. e., $p = 3$

does not form an exception. The existence of G_8 when $p > 3$ may be proved in exactly the same manner as when $p = 3$. Each of the five groups G_1, G_3, G_5, G_7, G_8 is conformal with the abelian group of type $(m-3, 1, 1)$, G_2 and G_4 are conformal with the abelian group of type $(m-3, 2)$ while G_6 is conformal with the one of type $(m-1, 1)$. Four of these groups (G_1, G_2, G_5, G_6) contain invariant cyclic subgroups of order p^{m-2} while these subgroups are conjugate, in sets of p , in the remaining four groups.

W. F. OSGOOD: *On a fundamental theorem...*

P. 278, l. 5. *After* point *insert* and no two curves corresponding to two distinct values of a will intersect each other.

E. J. WILCZYNSKI: *Geometry of a simultaneous system*

P. 359, l. 10 up. *For* form $y = \lambda\eta, z = \mu\zeta$ *read* form (2).

L. E. DICKSON: *Theory of linear groups in an arbitrary field.*

P. 370, l. 5. *For* $T_{s,-1} \cdots T_{3,-1}$ *read* $T_{2,-1} \cdots T_{s,-1}$.

P. 372, l. 4 up. *In* $A'_{13}: Y'_{12} = -Y'_{23}$, “ Y_{23} .

P. 377, l. 15. *For* $\Sigma s'$ “ $\Sigma s'$.

P. 384, l. 9. “ $+ Y_{13}\eta_3$ “ $+ Y_{12}\eta_3$.

P. 388, l. 15. “ *subscript* $-\lambda\nu^{-1}$ “ $-\lambda\nu$.

P. 388, l. 8 up. “ $p^{6n}\Omega_1$ “ $(p^{6n} - 1)\Omega_1$.

P. 390, l. 7 up. “ ξ_1 “ η_1 .

Pp. 383-391. For the simplicity of the group H' in the excluded case of modulus 2, see the report in the BULLETIN, November, 1902, of the Ninth Summer Meeting of the Society at Evanston.

VOLUME 3

O. STOLZ: *Zur Erklärung der Bogenlänge....*

P. 31, l. 17. *For* $\sum_r f_r d_r$ *read* $\sum_r f_r \delta_r$.

P. 35, l. 13. “ κ “ Δ .

L. E. DICKSON: *The groups of Steiner in problems of contact.*

P. 44, l. 22. *For* $(00 x_2 y_2 x_3 x_3 \cdots)$ *read* $(00 x_2 y_2 x_3 y_3 \cdots)$.

C. N. HASKINS: *On the invariants*

- P. 73, (2). For $\frac{\partial \xi_n}{\partial x_k}$ read $\frac{\partial \xi_r}{\partial x_k}$.
- P. 75, (10). " $a_{ik} \frac{\xi_r}{r_{i_1} i_2}$ " $a_{ik} \frac{\xi_r}{r_{i_1} i_2}$.
- P. 77, (1). " $a_{kr} \frac{\xi_r}{u_1 \dots i_\mu i_{\mu+1}}$ " $a_{kr} \frac{\xi_r}{u_1 \dots i_\mu i_{\mu+1}}$.
- P. 77, (2). " $a_{kr} \frac{\xi_r}{u_1 \dots i_\mu i_{\mu+1}}$ " $a_{kr} \frac{\xi_r}{u_1 \dots i_\mu i_{\mu+1}}$.
- P. 80. In equation (1) the left member is $(\lambda\mu, \nu\rho)$.
- P. 82, V. For ξ_n read ξ_r .
- P. 83, (9). " ξ_1 " $0\xi_1$.
- P. 86. The numerator of the fraction in the last line is $|C_{11} C_{22} C_{33}|$.

P. 87. For $\begin{vmatrix} a_{12} & 2Z & 3Z_1 & 2Z_2 \\ a_{22} & 0 & z_2 & 0 \\ a_{11} & 0 & 0 & z_1 \\ a_{21} & 2Z & 2Z_1 & 3Z_2 \end{vmatrix}$ read $\begin{vmatrix} a_{12} & 2Z & 3Z_1 & 2Z_3 \\ a_{22} & 0 & Z_2 & 0 \\ a_{11} & 0 & 0 & Z_1 \\ a_{21} & 2Z & 2Z_1 & 3Z_2 \end{vmatrix}$

E. B. VAN VLECK: *A determination of the number*

- P. 130, l. 16. This line should be $E\left(\frac{X+2}{2}\right) + E\left(\frac{Y+2}{2}\right) - 2$.
- P. 130, l. 17. For (22) read (21).
- P. 130, l. 10 up. " $\lambda - \lambda'_i - \lambda'_j + 1$ " $\lambda'_i + \lambda'_j > \lambda'_j + 1$.

E. H. MOORE: *On the projective axioms of geometry*.

- Pp. 142-158. In Hilbert's system I, II the axiom I4 is not redundant. The error in my proof of the dependence of I4 lies in the omission of a citation of I4 in proof of the second statement of ll. 21-24, p. 143. That statement, call it I4*, is in effect: If two planes have their respective sets of incident points identical, they are themselves identical. The axiom I4 does depend upon I1-3, 5, 7, II 1-3, 5 and I4*.—E. H. M.

- P. 143, l. 22. Insert the references (I2; I4).—The connection is in a 1-1 way.
- P. 146, l. 6 up. Omit indeed.
- P. 147, l. 4. For thus read hence.
- P. 155, l. 13. " segment " line.
- " α " α' .
- P. 156, l. 15. " k -space " $(k-1)$ -space.